

POSTER PRESENTATION

Open Access

# Assessment of the time course of chronic inflammation in the murine house dust mite model

Stephen M Jordan<sup>1\*</sup>, Robert Carrington<sup>1</sup>, David Lanham<sup>2</sup>, James Cartwright<sup>3</sup>, Rachel A Armstrong<sup>1</sup>, Kenneth G Meecham<sup>1</sup>

From 2nd Cross Company Respiratory Symposium  
Horsham, UK. 6-7 September 2012

House dust mite (HDM) allergens are associated with allergic disorders and the use of this clinically relevant allergen is increasing in animal models. We assessed the chronic inflammatory time course and the anti-inflammatory efficacy of a phosphodiesterase 4 inhibitor and a corticosteroid. BALB/c mice were challenged intranasally with HDM for 5 days/week for 5 weeks. Animals were sacrificed weekly 24 hours after final challenge and recruitment of inflammatory cells assessed in bronchoalveolar lavage fluid (BALF). Lung tissue was stained for the evaluation of a histopathological response. Roflumilast (10 mg/kg) and prednisolone (10 mg/kg) were administered orally twice daily from Week 3. Chronic HDM extract exposure resulted in significant recruitment of eosinophils, neutrophils, lymphocytes and macrophages as early as Week 1, peaking ( $1.13 \pm 0.32$ ,  $0.31 \pm 0.05$ ,  $0.66 \pm 0.10$  and  $0.33 \pm 0.04 \times 10^6$  cells/animal respectively) between Weeks 3 and 5. Within the lymphocyte population the proportion of B cells increased from 4 to 46% over the 5 week exposure period. Mice developed perivascular, peribronchiolar and alveolar inflammation which increased in severity during the five week exposure period. The inflammation was accompanied by epithelial and mucus cell hypertrophy/hyperplasia in the bronchi and bronchioles which reached maximum severity during Weeks 3 to 5. Perivascular/peribronchiolar fibrosis peaked in Week 5. Therapeutic treatment with prednisolone and roflumilast significantly ( $P < 0.001$ ) inhibited BALF cell recruitment and reduced the severity of the airway remodelling suggesting this model, in our laboratory, has the potential to test novel compounds for the treatment of allergic disorders.

#### Authors' details

<sup>1</sup>Department of Pharmacology, Huntingdon Life Sciences, Cambridgeshire, UK. <sup>2</sup>Department of Biomarkers, Bioanalysis and Clinical Sciences, Huntingdon Life Sciences, Cambridgeshire, UK. <sup>3</sup>Department of Pathology, Huntingdon Life Sciences, Cambridgeshire, UK.

Published: 14 August 2013

doi:10.1186/1476-9255-10-S1-P1

**Cite this article as:** Jordan et al.: Assessment of the time course of chronic inflammation in the murine house dust mite model. *Journal of Inflammation* 2013 **10**(Suppl 1):P1.

**Submit your next manuscript to BioMed Central  
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



<sup>1</sup>Department of Pharmacology, Huntingdon Life Sciences, Cambridgeshire, UK

Full list of author information is available at the end of the article